

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

UNCLASSIFIED

Contract No. W-7405-eng-48

Half-Life for Double Beta-Decay

C. A. Levine, A. Ghiorso, and G. T. Seaborg

October 21, 1949

Berkeley, California

<u>Standard Distribution</u>	<u>No. of Copies</u>
Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brookhaven National Laboratory	4
Bureau of Ships	1
Carbide and Carbon Chemicals Corporation (K-25 Plant)	4
Carbide and Carbon Chemicals Corporation (Y-12 Plant)	4
Columbia University (Failla)	1
Dow Chemical Company	1
General Electric Company, Richland	6
Hanford Operations Office	1
Iowa State College	1
Kneels Atomic Power Laboratory	4
Los Alamos	3
Massachusetts Institute of Technology (Gaudin)	1
Massachusetts Institute of Technology (Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	1
National Bureau of Standards	2
Naval Radiological Defense Laboratory	1
NEPA Project	1
New Brunswick Laboratory	1
New York Operations Office	2
North American Aviation, Inc.	1
Oak Ridge National Laboratory	12
Patent Branch, Washington	1
Sylvania Electric Products, Inc.	1
Technical Information Branch, ORE	15
UCLA Medical Research Laboratory (Warren)	1
University of California Radiation Laboratory	4
University of Rochester	2
Western Reserve University (Friedell)	1
Westinghouse Electric Corporation	2
Total	95


Radiation Laboratory
University of California
Berkeley, California

HALF-LIFE FOR DOUBLE BETA-DECAY

by C. A. Levine, A. Ghiorso, and G. T. Seaborg
Department of Chemistry and Radiation Laboratory
University of California, Berkeley, California

Fireman⁽¹⁾ has reported the results of a rather difficult beta-particle coincidence counting experiment in which the decay of Sn^{124} by the simultaneous emission of two negative beta-particles, with a half-life between 0.4×10^{16} years and 0.9×10^{16} years, seems to have been observed. This note reports the results obtained from a different and somewhat simpler method of looking for the phenomenon of simultaneous emission of two beta-particles. These results are negative so far and show that this process is considerably less probable in the case chosen by us than in that reported by Fireman.

Our method consists of looking in uranium samples for 90-year Pu^{238} which would come from U^{238} by the double beta-particle mechanism since Np^{238} is heavier than U^{238} , which in turn is substantially heavier than Pu^{238} , in the isobaric triplet $_{92}\text{U}^{238}$ - $_{93}\text{Np}^{238}$ - $_{94}\text{Pu}^{238}$. This chemical method of investigation is particularly applicable to this isobaric triplet because there appears to be no other mechanism to account for the Pu^{238} should it be found. The energetics of the situation are summarized in the following diagram, where the disintegration energies are derived from sources which may be traced through a recent compilation.⁽²⁾



The alpha-disintegration energy of Np^{238} is estimated from alpha-decay systematics.⁽³⁾

Our experiment consisted of taking 14 kg. of very pure, 6-year old, UO_3 and extracting and separating the plutonium fraction by chemical means. The method

consisted essentially of dissolving the oxide in nitric acid and precipitating Pu(IV) with lanthanum fluoride, followed by solution of the lanthanum fluoride and oxidation of the plutonium to Pu(VI), which was extracted into diethyl ether and then re-extracted into water. Similar cycles were repeated five times in order to separate completely from UX_1 and to reduce the amount of lanthanum carrier, after which the final sample was plated out on flat platinum with total carrier weight probably less than 50 micrograms. The use of tracer Pu^{239} in this separation established that the chemical yield amounted to 10%.

This final sample was measured for the presence of the 5.51 Mev alpha-particles of Pu^{238} on the alpha-pulse analyzer apparatus in this laboratory⁽⁴⁾. This analysis showed that the counting rate of the Pu^{238} α -particles at essentially 50% counting yield amounted to 0.00 ± 0.01 counts per minute above background. This indicates that the "half-life" of U^{238} for simultaneous emission of two beta-particles, for which a total energy of 1.1 Mev is available, is greater than 6×10^{18} years.

This experiment could be extended to reach longer half-lives through the use of larger and older sources of uranium such as pitchblende ore. In this case, of course, the plutonium fraction so isolated will contain a certain amount of Pu^{239} as has already been demonstrated⁽⁵⁾. The extraction of plutonium from a ton of pitchblende (50% uranium) with 10% yield could detect a half-life as long as some 10^{22} years for the same limits of counting accuracy.

This result appears to disagree with that of Fireman although it may not be possible to be positive about this in view of the difference in energies and atomic numbers and possible difference in degree of forbiddenness involved. The theory for the double beta-decay process sets widely differing ranges of half-life depending upon whether the process can take place without neutrino emission⁽⁶⁾. Fireman's results are in the range predicted for double beta-decay without neutrino emission while our half-life limit seems to be above this predicted range and perhaps points

toward the emission of two neutrinos in this process.

It is a pleasure to acknowledge the assistance of Dr. L. B. Magnusson in the chemical procedure. This work was performed under the auspices of the U. S. Atomic Energy Commission.

References

- (1) E. L. Fireman, Phys. Rev. 75, 323 (1949).
- (2) G. T. Seaborg and I. Perlman, Revs. Mod. Phys., 20, 585 (1948).
- (3) I. Perlman, A. Ghiorso, and G. T. Seaborg, Phys. Rev. 74, 1730 (1948).
- (4) A. Ghiorso, A. H. Jaffey, H. P. Robinson, and B. Weissbourd, National Nuclear Energy Series, Plutonium Project Record Vol. 14B "The Transuranium Elements: Research Papers", Paper No. 16.8 (McGraw-Hill Book Co., Inc., New York, 1949).
- (5) G. T. Seaborg and M. L. Perlman, J. Am. Chem. Soc. 70, 1571 (1948); National Nuclear Energy Series, Plutonium Project Record Vol. 14B "The Transuranium Elements: Research Papers", Paper No. 1.3 (McGraw-Hill Book Co., Inc., New York, 1949).
- (6) W. H. Furry, Phys. Rev. 56, 1184 (1939).

October 21, 1949